

Chapter 06 Anti-Derivatives

1) The anti derivative of the function $f(y) = \sec y \tan y$ is

- A) $\sec y \tan y$
- B) $\sec^2 y$
- C) $\sec y + c$
- D) $\tan y$

Answer: C

2) The anti derivative of zero is

- A) 1
- B) 0
- C) x
- D) constant

Answer: D

3) The anti derivative of the function $f(y) = 4^y$ is

- A) $4^{y+1} + C$
- B) $4^y \ln 4$
- C) $\frac{4^y}{\ln 4} + C$
- D) $4^y + C$

Answer: C

4) The anti derivative of the function $f(y) = \tan^2 y \cosec^2 y$ is

- A) $\tan^2 y + c$
- B) $\tan y + c$
- C) $\cosec^2 y + c$
- D) $\cosec y + c$

Answer: B

5) The anti derivative of the function $f(z) = (3z - 8)^5$ is

- A) $\frac{(3z-8)^6}{6} + c$
- B) $\frac{(3z-8)^6}{12} + c$
- C) $\frac{(3z-8)^6}{18} + c$
- D) $6(3z-8)^6 + c$

Answer: C

6) The value of $\int \frac{4y^3 - 4y}{y^4 - 2y^2 + 3} dy$ is

- A) $\ln(y^3 - 4y) + c$
- B) $\ln(y^4 - 2y^2 + 3) + c$
- C) $\ln(y^4 - 2y^2 + 3) + c$
- D) $\ln(4y^3 - 4y) + c$

Answer: C

7) The value of $\int \frac{1}{yl_n y} dy$ is

- A) $l_n |y| + c$
- B) $l_n \frac{1}{y} + c$
- C) $\frac{1}{y} + c$
- D) $l_n [l_n |y|] + c$

Answer: D

8) The value of $\int \frac{\sec^2 y}{\tan 45^\circ} dy$ is

- A) $l_n |\tan y| + c$
- B) $l_n |\tan 45^\circ| + c$
- C) $\tan y + c$
- D) $\sec y + c$

Answer: C

9) The anti derivative of the function $f(y) = y \tan 45^\circ$ is

- A) $\sec^2 45^\circ + C$
- B) $y \sec^2 45^\circ$
- C) $\frac{y^2}{2} \sec^2 45^\circ$
- D) $\frac{1}{2} y^2 + c$

Answer: D

10) The value of $\int (\tan 45^\circ) \sec^2 y dy$ is

- A) $\frac{(\tan 45^\circ)^4}{4} + c$
- B) $\frac{(\tan 45^\circ)^4}{4}$
- C) $\frac{(\tan 45^\circ)^4}{4} \tan y + c$
- D) $\tan y + c$

Answer: D

11) The value of $\int (\cot 45^\circ)^5 \cosec^2 y dy$

- A) $\frac{(\cot 45^\circ)^5}{6}$
- B) $-\frac{(\cot 45^\circ)^6}{6} + c$
- C) $\frac{(\cot 45^\circ)^6}{6} + c$

D) $-\cot y + c$

Answer: D

12) The value of $\int (\sin \alpha)^3 \cos x dx$ is

A) $\frac{(\sin \alpha)^4}{4}$

B) $\frac{(\sin \alpha)^4}{4} + c$

C) $\frac{(\sin \alpha)^3}{3} + c$

D) $(\sin \alpha)^3 \sin x + c$

Answer: D

13) The value of $\int \sin x \cos \alpha dx$

A) $\frac{(\sin x)^2}{2}$

B) $\frac{(\sin x)^2}{2} + c$

C) $-\cos x \cos \alpha + c$

D) $\cos x \cos \alpha + c$

Answer: C

14) the value of $\int \sec^3 \alpha dy$ is

A) $\tan \alpha + c$

B) $-\tan \alpha + c$

C) $y \sec^2 \alpha + c$

D) $y \tan \alpha + c$

Answer: C

15) The value of $\int \cos ec^2 a dx$ is

A) $-\cot a + c$

B) $\cot a + c$

C) $x \cot a + c$

D) $x \operatorname{cosec}^2 a + c$

Answer: D

16) The value of $\int \frac{\cot x \cot 45^\circ}{l_n \sin x} dx$ is

A) $l_n \sin x + c$

B) $l_n \cot x + c$

C) $l_n (l_n \sin x) + c$

D) $-l_n (l_n \cot x) + c$

Answer: C

17) The value of $\int \tan^2 y \sec^2 y dy$ is

A) $\frac{(\tan y)^3}{3} + c$

B) $\frac{(\tan y)^2}{2} + c$

C) $\frac{(\tan y)^3}{4} + c$

D) $\frac{1}{2}(\tan y)^3 + c$

Answer: A

18) The value of $\int (y^2 + y + 5)^4 (2y+1) dy$

A) $\frac{y^3}{3} + \frac{y^2}{2} + 5y + c$

B) $\frac{(y^2 + y + 5)^4}{4} (y^2 + y) + c$

C) $\frac{(2y+1)^5}{5} + c$

D) $\frac{(y^2 + y + 5)^5}{5} + c$

Answer: D

19) The partial fraction of $\frac{2x-1}{x(x-1)}$ are

A) $\frac{Ax+b}{x^2-x}$

B) $\frac{A}{x} + \frac{B}{x-1}$

C) $\frac{A}{x} - \frac{B}{x-1}$

D) $\frac{A}{x-1} - \frac{B}{x}$

Answer: B

20) In the form of partial fractions the rational function

$\frac{(3x^2-1)(2x+1)}{(x-1)(x^2+1)}$ can be written as

A) $\frac{A}{x-1} + \frac{B}{x^2+1}$

B) $\frac{A}{x-1} - \frac{B}{x^2+1}$

C) $\frac{A}{x-1} + \frac{Bx}{x^2+1}$

D) $\frac{A}{x-1} + \frac{Bx+c}{x^2+1}$

Answer: D

- 21) In the form of partial fractions the rational function $\frac{x^3 + x^2 + 2x + 3}{(x^2 + 1)(x^2 + 2)}$ can be written as

A) $\frac{A}{x^2 + 1} + \frac{B}{x^2 + 2}$
B) $\frac{Ax + B}{x^2 + 1} + \frac{Cx + D}{x^2 + 2}$
C) $\frac{Ax - B}{x^2 + 1} + \frac{Cx - D}{x^2 + 2}$
D) $\frac{A + B}{x^2 + 1} + \frac{C + D}{x^2 + 2}$

Answer: B

- 22) In the form of partial fraction the rational function $\frac{x^2 + 2x}{(x+1)^2(x^2 + 1)}$ can be written as

A) $\frac{A}{x+1} + \frac{B}{x^2 + 1}$
B) $\frac{A}{x+1} + \frac{Bx + C}{x^2 + 1}$
C) $\frac{A}{(x+1)^2} + \frac{Bx + C}{x^2 + 1}$
D) $\frac{A}{x+1} + \frac{B}{(x+1)^2} + \frac{Cx + D}{x^2 + 1}$

Answer: D

- 23) The value of $\int \frac{dx}{(x+2)^2 + 4}$ is

A) $\tan^{-1}(x+2)$
B) $\tan^{-1} \frac{(x+2)}{2}$
C) $\frac{1}{2} \tan^{-1} \frac{(x+2)}{2} + C$
D) $\frac{1}{4} \tan^{-1} \left(\frac{x+2}{4} \right) + C$

Answer: C

- 24) The value of $\int \frac{dx}{x^2 - 1}$ is

A) $\ln \left| \frac{x-1}{x+1} \right| + C$

B) $\ln \left| \frac{x+1}{x-1} \right| + C$

C) $\frac{1}{2} \ln \left| \frac{x+1}{x-1} \right| + C$

D) $\frac{1}{2} \ln \left| \frac{x-1}{x+1} \right| + C$

Answer: D

- 25) The value of $\int \frac{dx}{4-x^2}$ is

A) $\ln \left| \frac{2-x}{2+x} \right| + C$
B) $\ln \left| \frac{2+x}{2-x} \right| + C$
C) $\frac{1}{4} \ln \left| \frac{2+x}{2-x} \right| + C$
D) $\frac{1}{4} \ln \left| \frac{2-x}{2+x} \right| + C$

Answer: C

- 26) The value of $\int \frac{1+\cos x}{x+\sin x} dx$ is

A) $\ln |x+\sin x| + C$
B) $x + \sin x + C$
C) $\ln |x+\cos x| + C$
D) $\ln |\sin x + \cos x| + C$

Answer: A

- 27) The indefinite integral of the function $f(y) = \frac{2y}{y^2 + 1}$ is

A) $y^3 + C$
B) $\frac{y^3}{3} + y$
C) $y^2 + C$
D) $\ln |y^2 + 1| + C$

Answer: D

7. Integration by Substitution

- 1) The indefinite integral of $f(y) = \frac{e^y}{1+e^{2y}}$ is

A) $\ln |1+e^{2y}| + C$
B) $e^y + C$

C) $\tan^{-1} e^y + C$

D) $\cot^{-1} e^y + C$

Answer: C

2) The indefinite integral of $f(x) = \ln x$ is

A) $\frac{1}{x} + C$

B) $\frac{\ln x}{x} + C$

C) $\ln x - x + C$

D) $x \ln x - x + C$

Answer: D

3) The anti derivative of $f(y) = e^y \left(\cos^{-1} y - \frac{1}{\sqrt{1-y^2}} \right)$ is

A) $e^y \sin^{-1} y + C$

B) $e^y \cos^{-1} y + C$

C) $-e^y \cos^{-1} y + C$

D) $-e^y \sin^{-1} y + C$

Answer: B

4) The value of $\int e^y \left(\sec^{-1} y + \frac{1}{y\sqrt{y^2-1}} \right) dy$ is

A) $e^y \operatorname{cosec}^{-1} y + C$

B) $-e^y \operatorname{cosec}^{-1} y + C$

C) $e^y \sec^{-1} y + C$

D) $-e^y \sec^{-1} y + C$

Answer: C

5) The anti derivative

of

$$f(x) = e^x \left(\operatorname{cosec}^{-1} x - \frac{1}{x\sqrt{x^2-1}} \right)$$

A) $e^x \sec^{-1} x + C$

B) $e^x \operatorname{cosec}^{-1} x + C$

C) $-e^x \sec^{-1} x + C$

D) $-e^x \operatorname{cosec}^{-1} x + C$

Answer: B

6) The value of $\int e^y (y^3 + 3y^2) dy$ is

A) $e^y \left(\frac{y^4}{4} + y^3 \right) + C$

B) $e^y (y^4 + y^3) + C$

C) $e^y y^3 + C$

D) $3e^y y^2 + C$

Answer: C

7) The anti derivative of $f(x) = e^x x^3 + 3e^x x^2$ is

A) $e^x \frac{x^4}{4} + e^x x^3 + C$

B) $e^x x^4 + 3e^x x^2$

C) $e^x \frac{x^4}{4} + 3e^x x^3 + C$

D) $e^x x^3 + C$

Answer: D

8) The value of $\int (e^z \cos z - e^z \sin z) dz$ is

A) $-e^z \sin z + C$

B) $e^z \cos z + C$

C) $-e^z \cos z + C$

D) $e^z \sin z + C$

Answer: B

9) The definite integral of $f(y) = e^y \cot y - e^y \operatorname{cosec}^2 y$ is

A) $e^y \operatorname{cosec}^2 y + C$

B) $-e^y \operatorname{cosec}^2 y + C$

C) $-e^y \tan y + C$

D) $e^y \cot y + C$

Answer: D

10) The anti derivative of $f(z) = e^z \sec z + e^z \sec z \tan z$ is

A) $e^z \operatorname{cosec} z + C$

B) $e^z \sec z + C$

C) $-e^z \sec z \tan z + C$

D) $e^z \sec z \tan z + C$

Answer: B

11) The value of $\int (e^y \operatorname{cosec} y - e^y \operatorname{cosec} y \cot y) dy$ is

A) $e^y \operatorname{cosec} y \cot y + C$

B) $-e^y \operatorname{cosec} y \cot y + C$

C) $e^y \operatorname{cosec} y + C$

D) $-e^y \operatorname{cosec} y + C$


Answer: C

12) The value of $\int_{\frac{\pi}{4}}^{\frac{\pi}{2}} (\sec^2 y + \csc^2 y) dy$ is

- A) 0
- B) $\tan \frac{\pi}{4}$
- C) 1
- D) $-\cot \frac{\pi}{4}$

Answer: A

13) The shaded area in the figure can be represented by



- A) $\int_0^a f(x) dx$
- B) $\int_0^b f(x) dx$
- C) $\int_b^a f(x) dx$
- D) $\int_a^b f(x) dx$

Answer: D

14) The value of $\int_3^3 (x^3 + 3x^2 + 2x + 1) dx$ is

- A) 27
- B) 54
- C) 52
- D) 0

Answer: D

15) The value of $\int_1^3 (z^5 + 4z^4 + z^3) dz$ is

- A) 1
- B) 243
- C) 324

D) $-\int_3^1 (z^5 + 4z^4 + z^3) dz$

Answer: D

16) The value of $\int_0^{\frac{\pi}{3}} \sin y dy$ is

- A) 0
- B) 1
- C) $\frac{\sqrt{3}}{2}$
- D) $-\int_{\frac{\pi}{3}}^0 \sin y dy$

Answer: D

17) The value of $\int_0^{\frac{\pi}{4}} \tan y \sec^2 y dy$ is

- A) 0
- B) 1
- C) -1
- D) $-\int_{\frac{\pi}{4}}^0 \tan y \sec^2 y dy$

Answer: D

18) The value of $\int_1^3 (x^2 + 2x)^5 dx + \int_3^5 (y^2 + 2y)^5 dy$ is

- A) $\int_1^5 (x^2 + 2x)^5 dx$
- B) $\int_3^5 (y^2 + 2y)^5 dy$
- C) $\int_1^3 (y^2 + 2y)^5 dy$
- D) $\int_1^5 (y^2 + 2y)^6 dy$

Answer: A

19) The value of $\int_0^2 (y^3 + 3y^2)^6 dy + \int_2^4 (z^3 + 3z^2)^6 dz$ is

- A) 204
- B) 364

- C) $\int_0^4 (z^3 + 3z^2)^6 dz$
D) $\frac{3}{4}$

Answer: C

- 20) If f and g are continuous functions and $\int_1^3 f(y) dy = 8, \int_3^7 f(z) dz = 9$ then the value of $\int_1^7 f(z) dz$ is

- A) 7
B) 1
C) 6
D) 17

Answer: D

- 21) If f and g are continuous functions on (a, b) , s.t. $\int_a^b f(x) dx = 12$ and $\int_a^b g(x) dx = 5$ then the value of $\int_a^b [f(x) - g(x)] dx$ is

- A) $7ab$
B) $7(a - b)$
C) $7(b - a)$
D) 7

Answer: D

- 22) If f and g are continuous functions on $(1, 5)$, such that $\int_1^5 f(y) dy = 5$ and $\int_1^5 g(y) dy = 3$ then the value of $\int_1^5 [f(y) + g(y)] dy$ is

- A) 6
B) 8
C) 4
D) 2

Answer: B

- 23) If $\int_2^5 f(x) dx = 5$ then the value of $\int_2^5 5f(y) dy$ is

- A) 5
B) 3
C) 25
D) 10

Answer: C

- 24) If $\int_1^3 f(x) dx = 4$ then the value of $\int_3^1 f(y) dy$ is

- A) 2
B) -2
C) 3
D) -4

Answer: D

- 25) The solution of the equation $\frac{dy}{dx} = 2x$ is

- A) $y = 2x$
B) $y = x^2 + c$
C) $y = 2x^2 + c$
D) $y = 3x^2 + c$

Answer: B

- 26) Let f be continuous on $(1, 7)$ and $\int_1^7 f(x) dx = 9$,

- $\int_1^7 f(y) dy = 19$ then the value of $\int_3^7 f(z) dz$ is

- A) 10
B) 28
C) 7
D) 4

Answer: A

- 27) The solution of $\frac{dy}{dx} = \frac{x}{y}$ is

- A) $x^2 + c$
B) $y^2 = x^2 + c$
C) $y^2 = x + 2c$
D) $y^2 = 2x$

Answer: B

- 28) If the acceleration of a particle is given by $z = 2t$, then its velocity at any time t is:

- A) $2t^2 + c$
B) $3t^2 + c$
C) $t^2 + c$
D) 2

Answer: C

- 29) If the velocity of a particle moving in a straight line is given by $v = 3t^2$ then the distance traveled by it in the first T seconds is

- A) $3t^2 + c$
B) $t^3 + c$
C) $3t^2 + c$
D) $T^3 + c$

Answer: D

- 30) The solution of $\frac{dy}{dx} = \frac{1}{x}$ is

- A) $y = x^2 + c$

B) $y = l_n |Cx|$

C) $y = l_n \left| \frac{1}{x} \right|$

D) $y = \frac{1}{x} + c$

Answer: B